

PHY 211 – Exam 2 (Version 1)

Name (please print): _____

SUID: _____

Please circle your TA's name: Richard Prashant Raghav Francesco

It is very important that you print your name at the top of the exam page. Please do it before you read any questions!

Document your work. Use the back of each sheet if you run out of space.

1.[25 pts total] Mike throws a ball upward and toward the east at a 63° angle with the level ground and a speed of 22 m/s. Nancy drives east past Mike at 30 m/s at the instant he releases the ball.


a. [6 pts] What are the x and y components of the initial velocity of the ball in Mike's reference frame?

b. [3 pts] Is the ball heading east or west in Nancy's reference frame?

c. [8 pts] What is the ball's initial speed in Nancy's reference frame?

d. [8 pts] What is the ball's initial angle in Nancy's reference frame?

2. [25 pts total] The following diagram shows a 75 kg person holding onto a rope running through a pulley with a 100 kg block attached to it. Assume that the string and pulley are massless and that there is no friction in the pulley. The system is at rest.



- a. [5 pts] Draw the free-body diagrams for the person and for the block, labeling all of the forces acting on each. Make sure to label which free-body diagram is which.
- b. [10 pts] What is the tension in the rope?
- c. [10 pts] The block rests on an industrial-size scale. What is the reading of the scale (in Newtons)?

3. [25 pts total] A 0.050 kg wooden block at the bottom of a wooden ramp is given an initial speed up the ramp of 3.0 m/s. The ramp is inclined at 23° above level ground. The coefficient of kinetic friction of wood on wood is 0.20 and the coefficient of static friction of wood on wood is 0.50.

- [4pts] Draw a diagram of the situation and draw the free body diagram for the block.
- [9pts] How far up the ramp is the block when its speed slows to zero?
- [8pts] What is the maximum angle to which the ramp can be raised before the block placed at rest on the ramp will start to slide down? Justify your answer from Newton's Laws and our model of friction.
- [4pts] When the block's speed slows to zero in the above example does it start to slide back down, or does it stay where it is? Explain.

4. [25 pts total] A computer hard disk 8.0 cm in diameter is initially at rest. A small dot is painted on the edge of the disk. The disk accelerates at 600 rad/s^2 for $\frac{1}{2} \text{ s}$, then coasts at a steady angular velocity for another $\frac{1}{2} \text{ s}$.

a. [6 pts] What is the speed of the dot after $\frac{1}{2} \text{ s}$?

b. [6 pts] Through how many revolutions has the dot turned after $\frac{1}{2} \text{ s}$?

c. [6 pts] What is the speed of the dot after 1 s?

d. [7 pts] Through how many revolutions has it turned after 1 s?

Formula Sheet for Physics 211

$$\vec{v} = \frac{d\vec{x}}{dt}; \quad \vec{a} = \frac{d\vec{v}}{dt}$$

$$\Delta x = x(t_2) - x(t_1) = \text{signed area under the } v(t) \text{ curve from } t_1 \text{ to } t_2 = \int_{t_1}^{t_2} v(t) dt$$

$$\Delta v = v(t_2) - v(t_1) = \text{signed area under the } a(t) \text{ curve from } t_1 \text{ to } t_2 = \int_{t_1}^{t_2} a(t) dt$$

$$v_x = v_{0x} + a_x t; \quad x = x_0 + v_{0x} t + \frac{1}{2} a_x t^2; \quad v_x^2 = v_{0x}^2 + 2a_x \Delta x$$

$$\theta = \frac{s}{r} : \theta \text{ in radians}; \quad \omega = \frac{d\theta}{dt} \quad \alpha = \frac{d\omega}{dt}$$

$$a_c = a_{rad} = \frac{v^2}{r}; \quad T = \frac{2\pi r}{v}$$

$$v = \omega r : \omega \text{ in radians per unit time}; \quad a_{tan} = \alpha r : \alpha \text{ in radians per unit time squared}$$

$$\omega = \omega_0 + \alpha t; \quad \theta = \theta_0 + \omega_0 t + \frac{1}{2} \alpha t^2; \quad \omega^2 = \omega_0^2 + 2\alpha \Delta \theta$$

$$\vec{v} = v_x \hat{i} + v_y \hat{j}; \quad v_x = |v| \cos(\theta); \quad v_y = |v| \sin(\theta)$$

$$v = |\vec{v}| = \sqrt{v_x^2 + v_y^2}; \quad \theta = \tan^{-1}(v_y/v_x)$$

$$\vec{v}_{CB} = \vec{v}_{CA} + \vec{v}_{AB}$$

$$g = 9.8 m/s^2$$

$$\vec{F}_{net} = \sum_i \vec{F}_i = m\vec{a}; \quad \vec{F}_{AB} = -\vec{F}_{BA}$$

$$|\vec{F}_{fk}| = \mu_k N; \quad |\vec{F}_{fs}| \leq \mu_s N$$

$$|F_{net, radial}| = \frac{mv^2}{R}$$